

Pick up at Pharmacy  
over the counter:  
4-Dulcolax laxative  
tablets  
1-Large bottle of  
MiraLax

Your prep will be faxed to your  
Pharmacy\_\_\_\_\_

Name of Pharmacy

## CENTER FOR DIGESTIVE HEALTH

Gastroenterology Associates, P.C.

2222 53rd Avenue, Bettendorf, IA 52722 (563) 383-2686

William Davidson III, MD  
Richard L. Weyman IV, MD  
Young B. Huh, MD  
Jae H. Kim, MD

Day \_\_\_\_\_  
Date \_\_\_\_\_  
Arrival Time \_\_\_\_\_  
Procedure Time \_\_\_\_\_

### MIRALAX PREP

#### Special instructions:

The laxatives used for this procedure can be purchased over the counter at a drug or grocery store without a prescription. When purchasing Dulcolax (generic Bisacodyl) make sure that it is the LAXATIVE, not the stool softener.

#### One (1) WEEK before your exam:

- No popcorn, flax seed, or Olestra (fat substitute found in foods like potato chips) one week prior to your procedure.
- NO IRON within 3 days to your procedure. Example: no Multi-vitamin or iron supplement.
- If you tend to be constipated, take two tablespoons of Milk of Magnesia two days prior to your procedure.

#### One (1) DAY before your exam:

- **No solid foods are allowed!** You may have only clear liquids. See back side of sheet for clear liquid diet.
- Drink eight (8) oz of liquid every 1 to 2 hours when on the clear liquid diet while awake. This will help maintain hydration.

#### FIRST ROUND PREP INSTRUCTIONS:

Mix eight (8) tablespoons of MiraLax with thirty-two (32) ounces of clear liquid.

- Take two (2) Dulcolax at **12:00 PM**
- Drink an entire 8 ounce glass every 15 minutes.  
\_\_\_ 12:00 PM, \_\_\_ 12:15 PM, \_\_\_ 12:30 PM, \_\_\_ 12:45 PM

#### SECOND ROUND PREP INSTRUCTIONS:

Mix eight (8) tablespoons of MiraLax with thirty-two (32) ounces of clear liquid.

- Take two (2) Dulcolax at **3:00 PM**
- Drink an entire 8 ounce glass every 15 minutes.  
\_\_\_ 3:00 PM, \_\_\_ 3:15 PM, \_\_\_ 3:30 PM, \_\_\_ 3:45 PM \_\_\_

#### THIRD ROUND PREP INSTRUCTIONS:

Mix eight (8) tablespoons of MiraLax with thirty-two (32) ounces of clear liquid.

- Drink an entire 8 ounce glass every 15 minutes.  
\_\_\_ 7:00 PM, \_\_\_ 7:15PM, \_\_\_ 7:30 PM, \_\_\_ 7:45PM

#### Six (6) hours before your exam:

Date/Time: \_\_\_\_\_

- **Discontinue all clear liquids.**

**YOU MUST BE ACCOMPANIED BY A DRIVER WHO REMAINS AT THE CENTER FOR DIGESTIVE  
HEALTH DURING YOUR PROCEDURE**

**CENTER FOR DIGESTIVE HEALTH**

**Gastroenterology Associates, P.C.**  
2222 53<sup>rd</sup> Avenue, Bettendorf, IA 52722 (563) 383-2686

PLEASE CALL IF YOU ARE HAVE QUESTIONS, CONCERNS OR PROBLEMS  
WITH THE PREPARATION OR INSTRUCTIONS (563) 383-2686

**Common questions:**

- Bowel movements often occur about one hour after the first glass. They will continue periodically for approximately 1-2 hours after you finish drinking the last glass.
- A feeling of bloating or nausea may occur due to the large volume of liquid; this is usually temporary and will disappear once bowel movements begin.
- A small amount of Vaseline applied to the rectal area may help relieve any discomfort
- For morning procedures, on the day of the procedure if you are thirsty, drink only a few ounces of water **up to six hours prior** to your procedure time.
- Color of the stool should be yellow or clear and liquid. (Mountain Dew color is common.)

**CLEAR LIQUID DIET**  
**DO NOT DRINK ANYTHING RED OR PURPLE**

**CAN HAVE**

Apple Juice      Gatorade  
Tang              Popsicles  
HI-C Fruit Juice    Pop (diet or regular)  
Coffee              Water  
Broth or Bouillon    Tea  
Plain Jell-O (no fruit or sugar free)

**CANNOT HAVE**

Milk or Milk Products  
Orange Juice with Pulp  
Tomato Juice  
Prune Juice  
Alcohol

**YOU MUST BE ACCOMPANIED BY A DRIVER WHO REMAINS AT THE CENTER FOR DIGESTIVE  
HEALTH DURING YOUR PROCEDURE**